



PROFESSIONAL PHARMACY

1555 S. Pilgrim St. Moses Lake WA,
98837

Phone 1-509-764-2314
info@MLRX.com

Testosterone Pump

Before Using this Medicine:

“Inform your Doctor or Pharmacist if you are allergic to any prescription or non-prescription medicine; taking other medicines or have other medical problems; pregnant, planning a pregnancy, or breastfeeding. “Check with your Doctor, Pharmacists, or Nurse if information in this leaflet causes you special concern or if you want additional information about your medicine and its use. “Inform your doctor if you have any of these conditions: DIABETES; LIVER PROBLEMS; KIDNEY DISEASE.

Common use for this Medicine:

1- For the treatment of MALE HYPOGONADISM / TESTICULAR HYPOFUNCTION
2- For the treatment of OTHER ANTERIOR PITUITARY DISORDERS
This medication is used *for* low testosterone levels.

How to use this Medicine:

Use this medicine exactly as directed on the label, unless instructed differently by your doctor
.. This medication is for EXTERNAL USE only.
“WASH hands immediately after using medication.
.. Apply to clean dry skin, NOT to the scrotum.
Do not shower or bathe within 6 hours after application.
- Cover application site with clothing AFTER gel has dried.
.. Patient should read specific "PATIENT INFORMATION~ provided with this medicine before starting treatment and each time their prescription is renewed.

What to do if you miss a dose:

If you miss a dose, apply as soon as possible when remembered; and then resume regular dosing schedule.

How to store this medicine:

Keep in original closed container in a dark, cool, and dry place away from children. Discard unneeded medicine.

Call your doctor immediately if any of the following occurs:

Unusual, painful & prolonged erection; Enlargement of the breasts in males; Skin rash; Swelling of feet or lower legs; Muscle weakness; Depression; Muscle pain; Absence of breathing during sleep.

Other side effects:

Baldness; Abnormal hair growth; increased sensitivity to sunlight; Headache; Acne; Voice alteration; High blood pressure. If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

PRECAUTIONS AND WARNINGS'

"This medicine is CONTRAINDICATED for use during pregnancy. CALL YOUR DOCTOR IMMEDIATELY IF PREGNANT or intend to become pregnant.

.. This medication is harmful in pregnancy. Women of child bearing age should not come in contact with this medicine.

.. Notify doctor if you are pregnant, if pregnancy is suspected, or if you intend to become pregnant.

.. Male patients with prolonged, inappropriate, and painful erections should IMMEDIATELY DISCONTINUE the drug and consult a doctor or go to EMERGENCY.

.. Contact your doctor if any nausea, vomiting, changes in skin color, or ankle swelling occurs.

.. It is important that you keep your follow-up appointments with your doctor.

For any questions, either talk to your provider or your pharmacist, and a dose adjustment can be made.

For any questions, either talk to your healthcare provider or your pharmacist.

For any questions, either talk to your provider or your pharmacist, and a dose adjustment can be made.